

**ingentaconnect**

# Improvement of phytase thermostability by using sorghum liquor wastes supplemented with starch

**Authors:** Chen C-C.<sup>1</sup>; Hunag C-T.<sup>1</sup>; Cheng K-J.<sup>2</sup>

**Source:** Biotechnology Letters, Volume 23, Number 5, March 2001, pp. 331-333(3)

**Publisher:** Kluwer Academic Publishers

[next article > View Table of Contents](#)

[mark item](#)

[full text options](#)

## Abstract:

When a phytase solution, soluble starch, and sorghum liquor wastes were mixed at the ratio of 1:1:10 (v/w/w), the residual phytase activities after 30 min of treatment at 70 and 80 °C were respectively, about 90% and 18% of that at 37 °C. After 10 min treatment, the residual activity was 67% at 80 °C and 10% at 90 °C.

**Keywords:** phytase; sorghum liquor wastes; starch; thermostability

**Language:** English

**Document Type:** Regular paper

**Affiliations:** **1:** Department of Agricultural Chemistry, National Taiwan University, 1, Section 4, Roosevelt Road, Taipei 10617, Taiwan, ROC **2:** Institute of BioAgricultural Sciences, Academia Sinica, Taipei 11529, Taiwan, ROC

The full text article is available for purchase

**\$40.00 plus tax**

The exact price (including tax) will be displayed in your shopping cart before you check out. You will be able to remove this item from your shopping cart at any time before you have completed check-out.

**Add to cart**

[next article > View Table of Contents](#)

[Back to top](#)

© 2005 Ingenta [Terms and Conditions](#) | [Privacy Policy](#)

<http://www.ingentaconnect.com/content/klu/bile/2001/00000023/00000005/00323464;jsessionid=108ep5...> 8/22/2005